https://doi.org/10.1093/cercor/bhac468 Advance access publication date 17 December 2022 Original Article

Brain volumes, thicknesses, and surface areas as mediators of genetic factors and childhood adversity on intelligence

Camille M. Williams 1,*, Hugo Peyre^{1,2,3}, Franck Ramus¹

¹Laboratoire de Sciences Cognitives et Psycholinguistique, Département d'Études Cognitives, École Normale Supérieure, EHESS, CNRS, PSL University, 29 rue d'ulm, 75005, Paris, France,

²INSERM UMR 1141, Paris Diderot University, 48 Bd Sérurier, 75019, Paris, France,

³Department of Child and Adolescent Psychiatry, Robert Debré Hospital, APHP, 48 Bd Sérurier, 75019, Paris, France

*Corresponding author: LSCP, Département d'Études Cognitives, École Normale Supérieure, 29 rue d'Ulm, 75005 Paris, France. Email: williams.m.camille@gmail.com

Although genetic and environmental factors influence general intelligence (g-factor), few studies examined the neuroanatomical measures mediating environmental and genetic effects on intelligence. Here, we investigate the brain volumes, cortical mean thicknesses, and cortical surface areas mediating the effects of the g-factor polygenic score (gPGS) and childhood adversity on the g-factor in the UK Biobank. We first examined the global and regional brain measures that contribute to the g-factor. Most regions contributed to the g-factor through global brain size. Parieto-frontal integration theory (P-FIT) regions were not more associated with the g-factor than non-PFIT regions. After adjusting for global brain size and regional associations, only a few regions predicted intelligence and were included in the mediation analyses. We conducted mediation analyses on global measures, regional volumes, mean thicknesses, and surface areas, separately. Total brain volume mediated 7.04% of the gPGS' effect on the g-factor and 2.50% of childhood adversity's effects mediated by individual regional volumes, surfaces, and mean thicknesses was 10–15 times smaller. Therefore, genetic and environmental effects on intelligence may be mediated to a larger extent by other brain properties.

Key words: intelligence; cerebral measures; cerebellar volumes; polygenic score; childhood adversity.

Introduction

The positive correlation in performance across cognitive tests can be reduced to a single dimension: the general intelligence factor (g-factor), which reflects a person's general cognitive performance. A number of studies suggest that individual differences in brain structure are associated with individual differences in intelligence (for review Deary et al. 2010, 2021) and that genes and early environmental factors, such as childhood adversity, be responsible for interindividual variability in brain structure (Enlow et al. 2012; McGuire and Jackson 2020). Although several studies examined the genetic and neurological basis of intelligence separately, there are relatively few studies investigating how genetic, environmental, and neurological factors simultaneously influence intelligence due to the lack of sufficiently rich and large datasets (for review Deary et al. 2021). Thus, this paper aims to capitalize on the richness of the UK Biobank-a largescale prospective study with neural, genetic, environmental, and behavioral data-to identify the neuroanatomical measures (e.g. brain volumes) mediating the genetic and environmental effects on intelligence.

Intelligence is heritable, with genetic differences accounting for about 50% of the differences in intelligence (Haworth et al. 2010; Polderman et al. 2015). Genome-wide association studies (GWASs) identify genetic differences linked to variations in intelligence by pinpointing the single-nucleotide polymorphisms (SNPs) that contribute to differences in intelligence. SNPs that vary with intelligence scores are typically associated with brain-expressed genes (Johnson et al. 2016; Lee et al. 2018) that are linked to a range of neuronal classes and processes, such as synaptic and neuron differentiation (Hill et al. 2019). SNP variations are therefore thought to be associated with differences at the macroscopic cerebral level.

SNP variations related to a trait can be summarized into a single score: a polygenic score (PGS), which reflects an individual's genetic predisposition to a given phenotype. A PGS is derived from the sum of the effect allele at each SNP that is weighted by the SNP's effect on a trait (estimated in a GWAS). The PGS of cognitive performance (i.e. measured by a verbal numerical score in the UK Biobank and a g-factor in Cognitive Genomics and Cohorts for Heart and Aging Research in Genomic Epidemiology consortiums) predicted up to 10.6% of the variance in cognitive performance in an independent sample (Lee et al. 2018). Although PGSs do not currently explain enough variance in intelligence to accurately predict individual intelligence scores (Morris et al. 2020), PGSs are valuable measures of genetic factors at the population level. Educational attainment and cognitive performance PGSs are increasingly used to disentangle environmental from genetic effects on educational and life outcomes (e.g. Bates et al. 2018; Rimfeld et al. 2018; von Stumm et al. 2020; Saarentaus et al. 2021).

Since genetic and environmental effects act on intelligence via the brain, numerous studies investigated the neural correlates of intelligence. The most well-replicated association is the positive Downloaded from https://academic.oup.com/cercor/article/33/10/5885/6918725 by INIST-CNRS INEE INSB user on 01 June 2022

correlation of total brain volume (TBV) with intelligence scores, ranging from r = 0.24 to 0.31 (Pietschnig et al. 2015; Gignac and Bates 2017; Cox et al. 2019). Beyond overall brain size, the parieto-frontal integration theory (P-FIT; Jung and Haier 2007) is the most supported theory on the regional correlates of intelligence. Although studies report additional regional associations with the g-factor that go beyond those predicted by the P-FIT, structural (gray and white matter volumes), diffusion, and functional imaging studies find that intelligence scores are associated with the lateral and medial frontal, parietal, lateral temporal, and lateral occipital cortices and their underlying white matter connectivity (e.g. arcuate fasciculus; e.g. Cox et al. 2019; Deary et al. 2021; Gur et al. 2021).

Previous UK Biobank studies examined the brain correlates of intelligence (Cox et al. 2016, 2019). The authors reported consistent associations with the P-FIT theory, such as stronger associations in the frontal pole, and the paracingulate, as well as less consistent associations with the P-FIT theory, such as weak associations in inferior frontal and superior parietal areas. They also found associations in the insula and precuneus/posterior cingulate volumes (Cox et al. 2019), which were more recently implicated in general intelligence (Basten et al. 2015). As for subcortical volumes, the UK Biobank study found that the thalamic volumes were most associated with verbal numerical reasoning ($\beta = 0.23$). Finally, the authors reported that many of these regions still predicted intelligence when adjusting for TBV, suggesting that some regions make a unique contribution to intelligence that goes beyond TBV.

Although genetic and brain correlates of general intelligence (g-factor) have largely been studied, only 2 studies, to our knowledge, examined the extent to which neural measures mediate the effect of the g-factor polygenic score (gPGS) on the phenotypic g-factor. One study using vertex-wise mediation analyses of cortical thickness and cortical surface areas reported that the association between the gPGS and the phenotypic g-factor was mediated by the cortical thicknesses and surface areas of the anterior cingulate cortex, the prefrontal cortex, the insula, the medial temporal cortex, and inferior parietal cortex up to 0.75% in IMAGEN (N = 1,651) and 0.77% in IntegraMooDS (N = 742; Lett et al. 2020). In other words, these regions explained 20%-40% of the variance explained by the gPGS on the g-factor (3%-5%). A preprint on 550 adults, which used the same summary statistics of intelligence (Savage et al. 2018) as the above study to create their gPGSs, found that 2 intraparietal areas and the posterior temporal cortex surface areas mediated the effect of the gPGS on the g-factor (Gencet al. 2022). These mediation studies suggest that specific cortical regions mediate the effect of the gPGS on the phenotypic g-factor.

However, the extent to which additional regions, such as subcortical and cerebellar volumes, mediate the effect of the gPGS on intelligence has yet to be investigated. Moreover, the associations between cortical regions, genetics, and intelligence warrant further support and have not been investigated using the finergrained segmentations of the Destrieux atlas. Therefore, our first aim was to examine whether subcortical volumes, cerebellar volumes, cortical volumes, cortical thicknesses, and cortical surface areas mediate the effects of the gPGS and the phenotypic g-factor with a more predictive gPGS in the UK Biobank.

Since early adversity is associated with a decrease in intelligence and cognitive function later in life (Enlow et al. 2012; McGuire and Jackson 2020) and may have lasting biological and cerebral effects changes in childhood and adulthood (Lupien et al. 2009; Dye 2018), our second aim was to examine whether the regions that mediate the gPGS' effect on the g-factor also mediate childhood adversity's effect on the g-factor. Taken together, this paper contributes to our understanding of the neuroanatomical measures mediating genetic (gPGS) and early environmental (childhood adversity) effects on intelligence (g-factor).

Materials and methods

Analyses were run on R (R Core Team 2022) and preregistered here: https://osf.io/ec97u/?view_only=4b366bd7ed2442a1a9f64 bfcc2fe0946

Participants

The UK Biobank is a large prospective study with phenotypic, genotypic, and neuroimaging data from more than 500,000 participants. Participants were recruited between 2006 and 2010, from the vicinity of 22 assessment centers in England, Wales, and Scotland, with an age range for inclusion of 40–69 years. Data collection continues up to date.

All participants provided informed consent ("Resources tab" at https://biobank.ctsu.ox.ac.uk/crystal/field.cgi?id=200). The UK Biobank received ethical approval from the Research Ethics Committee (reference 11/NW/0382) and the present study was conducted based on application 46007.

We used the g-factor and the g-factor PGS derived in a previous paper (Williams et al. 2022). In brief, we included participants whose combination of cognitive tests allowed for a correlation with the complete g-factor of 0.70 or higher (N = 261,701). This threshold was used to maximize the robustness of the factor and the number of participants with a g-factor. The g-factor PGSs were created using sBayesR (Lloyd-Jones et al. 2019), from g-factor GWAS summary statistics that were obtained from UK Biobank participants that did not have neuroimaging data. In the present paper, we used the PGSs of individuals of British ancestry who did not have first or second-degree cousins, siblings, and parent offsprings in the discovery of GWAS. We residualized the gPGS for birth year and the first 40 principal components (PCs) prior to its inclusion in the mediation models.

In a previous paper (Williams et al. 2021), we analyzed the image-derived phenotypes from the first magnetic imaging resonance (MRI) visit generated by an image-processing pipeline developed and run by the UK Biobank Imaging team (Miller et al. 2016; Alfaro-Almagro et al. 2018) and reported 40,028 individuals with sex, age at MRI, and TBV data after excluding outliers. From here on, age at the first MRI visit will be referred to as age. From the 40,028 individuals with neuroimaging data, there were 39,131 participants with a g-factor of good quality (Table 1).

Image-derived phenotypes

We used the 10 global and 620 regional imaging phenotypes previously examined by Williams et al. (2021). The global phenotypes include TBV, total mean cortical thickness (MCT), total surface area (TSA), subcortical gray matter volume (GMV), cortical GMV, cerebral white matter volume (WMV), cerebellar GMV, cerebellar WMV, the brainstem volume, and cerebral spinal fluid (CSF), whereas the regional phenotypes include 444 cortical regions (148 volumes, 148 surface areas, and 148 cortical thicknesses) from the Freesurfer a2009s segmentations (Destrieux Atlas, datafield 197); 116 whole segmentations and subsegmentations of the amygdala, hippocampus, and thalamus and subsegmentations of the brainstem (Freesurfer subsegmentations, data-field 191); 28 cerebellum GMV segmentations from the FAST segmentations (data-field 1101); and 32 subcortical, white matter, and

Table 1. Descriptive statistics of the variables included in each analysis.

Analyses	Max N	Variable	Mean	SD	Median
Regressions	39,131	Age (years)	64.05	7.52	64.50
		g-factor	0.00	1.00	-0.01
Mediation models	28,917	Age (years)	64.04	7.46	64.50
with g-factor PGS		g-factor PGS	0.00	1.00	-0.05
		g-factor	0.20	1.05	0.18
Mediation models	19,956	Age (years)	64.08	7.42	64.58
with g-factor PGS and		g-factor PGS	0.03	0.99	0.02
childhood adversity		g-factor	0.27	1.03	0.26
-		- Childhood adversity	-0.03	1.39	-0.50

g-factor: general intelligence score. g-factor is scaled for the regressions. PGS: polygenic score. PGS is scaled before being residualized for birth year and the first 40 principal components of the genotyped data. Childhood adversity was scaled on the sample (N about 150,000) from which it was created. Max N: Maximum number of participants included in the analyses. Corresponds to N for TBV but will be reduced when including other brain measures when data for a region are missing.

Table 2.	PC loadings	of childhood	abuse and	stressor variables.
----------	-------------	--------------	-----------	---------------------

Variable	PC1	PC2	PC3	PC4	PC5
Felt hated by a family member as a child	0.528	-0.070	0.328	-0.114	0.772
Felt loved as a child	0.511	0.228	0.081	-0.682	-0.464
Someone to take to the doctor when needed as a child	0.351	0.620	-0.616	0.313	0.124
Physically abused by family as a child	0.491	-0.141	0.383	0.648	-0.416
Sexually molested as a child	0.310	-0.734	-0.600	-0.062	-0.033
Cumulative proportion of explained variance	0.419	0.599	0.772	0.896	1.000

PC: principal component of principal component analysis. Statements were rated on a 5 point scale from "never true" to "very often true". "Felt loved..." and "someone to take to the doctor..." we're reverse coded.

ventricle volumes from the Freesurfer ASEG segmentations (datafield 190). Freesurfer subcortical segmentations for the caudate, putamen, accumbens, and pallidum were used instead of the preregistered FIRST volumes, for segmentation consistency with the other subcortical and cortical volumes that were segmented from Freesurfer.

Although we did not preregister that we would examine the effects of the left and right measures of the whole thalamus, hippocampus, and amygdala because we focused on the association of their subsegmentations with the g-factor, we ran these exploratory analyses to facilitate result comparison with previous studies and examine whether associations at the subcortical subsegmentation level manifested at the global level.

Childhood adversity score

A childhood adversity score was created from questions in the UK Biobank on childhood abuse and social stressors. Childhood abuse was measured with data fields 20488 "When I was growing up... People in my family hit me so hard that it left me with bruises or marks") and 20490 ("When I was growing up... Someone molested me (sexually)") and childhood stressors were measured with data fields 20487 ("When I was growing up... I felt that someone in my family hated me"), 20489 ("When I was growing up... I felt loved"), and 20491 ("When I was growing up... There was someone to take me to the doctor if I needed it"). All questions are rated from 0 (Never True) to 4 (Very Often True). So that all indicators are in the same direction, we subtracted data field 20489 and data field 20491 responses from 4 (reverse coding). We conducted a PCA (prcomp function, R Core Team 2022) on the scores from these questions and extracted the first principal component (PC1) as our measure of childhood adversity, which captured 42% of the variance across questions (Table 2).

Statistical analyses

We refer to the phenotypic g-factor as the g-factor and the g-factor PGS as the gPGS. A residualized gPGS was created by adjusting the

gPGS for birth year and the first 40 principal components of the genotyped data and is referred to as the gPGS from here on out.

In the regression analyses, all continuous variables were meancentered and divided by 1 SD. Females were coded 0.5 and males— 0.5. Linear regressions were conducted using the *lm* function.

Mediation models were conducted within the SEM framework with *lavaan* (Rosseel 2012). For all mediation models described in the present study, indirect effects were calculated using the product method and we estimated all possible correlations between mediators (see Supplemental Tables D and E). We estimated direct paths (exposure to an outcome) and indirect paths (exposure to mediators to outcomes) and adjusted the mediators and outcome for age (linear and quadratic), sex, and their interactions in the *lavaan* framework. Sex was coded 0 for males and 1 for females. By default, correlations between exogenous variables (i.e. gPGS and childhood adversity) are fixed at their sample values in *lavaan*. We report the Monte Carlo CIs with 20,000 iterations for the mediation effects using the *semTools* package (Jorgensen et al. 2022) in Supplemental Tables D and E.

What global measures predict the phenotypic g-factor?

We first estimated the effect of TBV and the CSF on the phenotypic g-factor, while adjusting for sex, age (quadratic and linear), their interactions, and scanner site (Equation 1, where i refers to an individual).

$$\begin{split} g-factor_i &= Intercept + TBV_i + CSF_i + Sex_i + Age_i \\ &+ Age^2_i + Age^* \; Sex_i + Age^2 * Sex_i + e \end{split} \tag{1}$$

To identify the global measures driving the predictive effect of TBV on the g-factor, we simultaneously estimated the effect of total MCT, TSA, subcortical GMV cortical GMV, cerebral WMV, cerebellar GMV, cerebellar WMV, the brainstem volume, and CSF on the phenotypic g-factor, while adjusting for sex, age (quadratic and linear), their interactions, and scanner site (Equation 2, where i refers to an individual).

$$\begin{array}{l} g-factor_{i}=Intercept+Total Mean Cortical Thickness_{i}\\ +Total Surface Area_{i}+Cerebral WMV_{i}+Cerebellar GMV_{i}\\ +Cerebellar WMV_{i}+\\ Subcortical Volumes_{i}+Brainstem_{i}+CSF_{i}+Sex_{i}+Age_{i}\\ +Age^{2}_{i}+Age*Sex_{i}+Age^{2}*Sex_{i}+e \end{array}$$

$$(2)$$

We applied a Bonferroni correction to correct for multiple comparisons: we considered that a regional measure significantly predicts the g-factor when P < 0.05/N, N: the number of coefficients of interest, which was 2 for Equation (1) and 8 for Equation (2).

What global measures mediate the gPGS' effects on the g-factor?

We ran 2 mediation models using the sem function in the *lavaan* package (Rosseel 2012): one with the significant predictors of the g-factor from Equation (1) and one with those from Equation (2). The gPGS was the exposure, global volume(s) the mediator(s), and the g-factor served as the outcome.

We set a lenient P-value threshold to 0.05 and a stricter one to P < 0.05/N (N: the number of regional and global measures included in the model of interest) as a Bonferroni correction to correct for multiple comparisons. Good fit was established with a comparative fit index (CFI) > 0.95, a root mean square error approximation (RMSEA) < 0.06, and a standardized root mean square residual (SRMR) < 0.08 (Hu and Bentler 1999).

Do global measures mediate the gPGS' and childhood adversity's effects on the g-factor?

We applied the same mediation models as in What global measures mediate the gPGS' effects on the g-factor? section except that we added childhood adversity as additional exposure and estimated its direct and indirect paths through the global measure(s) to the g-factor. *Lavaan* considers correlations between predictors without estimating them. We set a lenient P-value threshold to 0.05 and a stricter one to P < 0.05/N (N: the number of regional and global measures included in the model of interest) as a Bonferroni correction to correct for multiple comparisons.

What regional measures predict the g-factor?

The aim was to identify the regions that contribute more to the g-factor than what is predicted given their size: This includes (i) regions that significantly predict the g-factor after adjusting for brain size and are positive with or without adjusting for global brain size and (ii) regions that significantly predict the g-factor after adjusting for brain size and are negative with or without adjusting for global brain size.

To do so, we ran Equations (3) and (4) (i refers to an individual, the regional measure N corresponds to a regional volume, thickness, or surface, and the global measure to TBV for volumes, total MCT for mean thicknesses, and TSA for surface areas). We applied a Bonferroni correction to correct for multiple comparisons: The significance threshold was set to P < 0.05/N (N: the number of coefficients of interest, which was 148 for surfaces, 148 for thicknesses, and 311 for volumes).

 $g - factor_i = Intercept + Regional Measure^{N_i} + Sex_i + Age_i$

 $+Age^{2}_{i} + Age * Sex_{i} + Age^{2} * Sex_{i} + e$ (3)

$$\begin{array}{l} g-factor_{i}=Intercept+Regional\ Measure^{N}{}_{i}\\ +Global\ Measure^{N}{}_{i}+Sex_{i}+Age_{i}+Age^{2}{}_{i}+Age\ast Sex_{i} \end{array}$$

 $+Age^2 * Sex_i + e$ (4)

Although this was not preregistered, we tested whether the P-FIT theory accurately predicted the neuroanatomical measures most associated with the g-factor. We mapped Brodmann areas from the P-FIT (Jung and Haier 2007; Colom et al. 2010; Haier and Jung 2018) to the regions of the Destrieux Atlas based on the region names and the description of their location. The 14 Brodmann areas that predict g according to the P-FIT were mapped to 62 of the 148 Destrieux segmentations. Several Destrieux regions were matched to the same Brodmann area and several Brodmann areas were matched to a single Destrieux region. Therefore, 12 Destrieux regions were matched twice to the Brodmann areas, yielding 74 P-FIT Destrieux regions out of 160 Destrieux regions (Supplemental Table F1)

We then examined (i) whether P-FIT regions had a larger effect on the g-factor than non-P-FIT regions, by comparing the distribution of effect sizes of the 2 sets of regions with a t-test; and (ii) whether P-FIT regions were overrepresented among the top 20 or 30 regions with the largest associations with the g-factor, using a chi-square test. We ran these analyses twice, for raw regional volumes, surfaces, and thicknesses, and for those adjusted on global brain measures. For the analysis of unadjusted regional volumes and surface areas, given that they were all significantly associated with the g-factor, we restricted the analysis to the regions showing the top N associations.

Do the regional measures that predict the g-factor independently from brain size still predict the g-factor when entered in the same model?

Based on Equations (3) and (4), we selected the regions that still significantly and positively or negatively predicted the g-factor after adjusting for brain size. Because brain regions are correlated, their effect on the g-factor may be shared across regions even if they are independent of global brain size. Therefore, to avoid redundancy, we examined whether these regions still predicted the g-factor when simultaneously entered into a regression model predicting the g-factor (Equation 5, where N refers to a region, i to an individual, and the global measure to TBV for volumes, total MCT for mean thicknesses, and TSA for surface areas). We applied a Bonferroni correction to correct for multiple comparisons: The significance threshold was set to P < 0.05/N (N: the number of regional measures included in the model of interest).

 $\begin{array}{l} g-factor_{i}=Intercept+Regional\ Measure^{^{1}}_{\ i}\\ +\ Regional\ Measure^{^{2}}_{\ i}\\ & \cdots\\ +Regional\ Measure^{^{N}}_{\ i}\\ +\ Global\ Measure^{^{N}}_{\ i}+Sex_{i}+Age_{i}+Age_{i}^{2}\\ +Age*Sex_{i}+Age^{^{2}}*Sex_{i}+e \end{array}$

Do regional measures mediate the gPGS' effects on the g-factor?

For volumes, thicknesses, and surface areas separately, we examined the simultaneous mediation of the global measure and the

(5)



Fig. 1. Meditating effect of global cerebral measures on the g-factor PGS's effect on the g-factor with and without including childhood adversity. g-factor, general intelligence factor; PGS, polygenic scores. Coefficients correspond to direct effects. Fit of TBV models: CFI=1.00, SRMR=0.00, RMSEA=0.00. Fit of global measures model: CFI=1.00, SRMR=0.01, RMSEA=0.03. The PGS is adjusted for birth year and the first 40 principal components of the genotyped data. Cerebral measures and the g-factor are adjusted for sex, age, age², age by sex, age² by sex, and scanner site. By default, correlations between exogenous variables are fixed at their sample values in *lavaan* (Rosseel 2012). For simplicity, the error variance of each variable and correlations between mediators are not specified in the figure (Supplemental Tables D and E).

significant regional measures from Equation (5) with the sem function from the *lavaan* package (Rosseel 2012). The gPGS was the exposure, global and regional measures were the mediators, and the g-factor served as the outcome. We set a lenient P-value threshold to 0.05 and a stricter one to P < 0.05/N (N: the number of regional and global measures included in the model of interest) as a Bonferroni correction to correct for multiple comparisons.

Do regional measures mediate the gPGS' and childhood adversity's effects on the g-factor?

We applied the same mediation models as in Do regional measures mediate the gPGS' effects on the g-factor? section except that we added childhood adversity as additional exposure and estimated its direct and indirect paths through regional and global measures to g-factor. We set a lenient *P*-value threshold to 0.05 and a stricter one to P < 0.05/N (N: the number of regional and global measures included in the model of interest) as a Bonferroni correction to correct for multiple comparisons.

Results What global measures predict the phenotypic g-factor?

Greater TBV was associated with a greater g-factor (β = 0.24, SE = 0.006, P = 6.99e-297) and CSF did not predict the g-factor (Supplemental Table B1). When dividing TBV into its subcomponents, we found that greater TSA (β = 0.14, SE = 0.012, P = 2.96e-31), total MCT (β = 0.04, SE = 0.006, P = 7.11e-12), cerebellar GMV (β = 0.08, SE = 0.008, P = 2.38e-26), and cerebral WMV (β = 0.05, SE = 0.013, P = 1.03e-04) were associated with an increase in the g-factor (Fig. 1; Supplemental Table B2). TBV explained 3.4% of the variance in the g-factor, whereas the global measures explained 3.6% of the variance in the g-factor.

What global measures mediate the gPGS' effect on the g-factor?

In the mediation model with TBV as the sole global mediator, TBV mediated 5.70% of the gPGS' effect on the g-factor (Supplemental Table D1).

In the mediation model with several global measures as mediators, TSA mediates 3.32%, cerebellar GMV 1.01%, and cerebral WMV 1.28% of the gPGS' effect on the g-factor (Fig. 1; Supplemental Table D2).

What global measures mediate the gPGS' and childhood adversity's effect on the g-factor?

In the mediation model with TBV as the sole global mediator, TBV mediated 7.03% of the gPGS' effect on the g-factor and mediated 2.49% of childhood adversity's effect on the g-factor (Supplemental Table E1).

In the mediation model with several global measures as mediators, TSA mediates 3.68%, cerebellar GMV 1.88%, and cerebral WMV 1.47% of the gPGS' effect on the g-factor and TSA mediates 1.19%, cerebellar GMV 0.56%, and cerebral WMV 0.96% of childhood adversity's effect on the g-factor (Fig. 1; Supplemental Table E2).

What regional measures predict the g-factor?

Regression results are available in Supplemental Table B (full models) and Supplemental Table C (regional estimates). Figure 2 shows the g-factor estimate by volume, surface area, or thickness estimate when including or excluding the global measure in the regression model (see Supplemental Files S8–S10 for brain maps). Figures of the estimate by region for each type of possible change in significance or estimate (Table 3) are available in Supplemental Files S1–3.

There were 242 out of 311 volumes, 130 out of 148 surface areas, and 6 out of 148 mean thicknesses, which were no longer significant after adjusting for global brain size (Table 3; Supplemental B3–4 and C1–3; Supplemental Files S1–3). We found that the size of volumes and surface areas was significantly and positively correlated with the magnitude of its association with the g-factor without adjusting for global measures and that this association was no longer significant after adjusting for global brain size (Supplemental File S4). Therefore, regional volumes and surfaces mainly contributed to the g-factor through global brain size.

There were 40 volumes (mainly cerebellar GMVs, subcortical thalamic and hippocampal nuclei, and a few cortical volumes), 12 cortical surface areas (mostly frontal), and 21 cortical mean thicknesses (mostly temporal; Table 3; Supplemental B5–6 and C1–3; Supplemental Files S1–3) that were significant and had positive estimates after adjusting for brain size, suggesting that



Model • With Global Measure • Without Global Measure

Fig. 2. Standardized estimate (Std Beta) of a region's association with the g-factor with (black) and without (light grey) adjusting for a global measure. Each point corresponds to a region. Region names are not shown for clarity. Global measure: TBV for volumes, total surface area for surface areas, and total mean cortical thickness for thicknesses.

Table 3. N	umber of volumes,	surface areas,	and mean	thicknesses	by types	of change ii	n significance	e and estimate b	etween mo	odels
without ar	nd with global brair	n size.				-				

	Volumes		Surface areas		Mean thicknesses	
Type of change between models	N	%	N	%	N	%
No longer significant	242	77.81	130	87.84	6	4.05
Still significant positive	40	12.86	12	8.11	21	14.19
Still significant negative	2	0.64	0	0.00	7	4.73
Still not significant	5	1.61	0	0.00	104	70.27
Still significant, positive to negative	18	5.79	6	4.05	0	0.00
Becomes significant and Negative	4	1.29	0	0.00	10	6.76
Total	311	100.00	148	100.00	148	100.00

Global brain size: total brain volume for volumes, total mean cortical thickness for thicknesses, and total surface areas for surface areas.

they contribute more (positively) to the g-factor than what is expected given their size.

There were 2 ventricular volumes and 7 cortical mean thicknesses (the left and right pericallosal sulci, the left and right anterior cingulate gyri and sulci, right occipital pole, left suborbital sulcus, and the right frontal marginal gyrus and sulcus; Table 3; Supplemental B5–6 and C1–3; Supplemental Files S1–3) that were significant and had negative estimates after adjusting for brain size, suggesting that they contribute more (negative) to the gfactor than what is expected given their size.

There were 18 volumes and 6 surface areas that were still significant but had their estimates switch from positive to negative, suggesting that they contribute less to the g-factor than what is expected given their contribution to global brain size. For volumes, these regions included the right and left caudate, the right and left lingual gyrus, left and right pericallosal sulci, the right posteriorventral part of the cingulate gyrus (isthmus), the right occipital pole, the right superior parietal gyrus, the hippocampal tail, and several subthalamic nuclei. As for surfaces, these regions included the left and right postcentral sulci, the left paracentral gyrus and sulcus, the middle anterior cingulate gyrus sand sulcus, the left lingual gyrus, and the left posterior-ventral part of the cingulate gyrus (isthmus).

Finally, there were 10 mean thicknesses, 3 ventricular volumes, and 1 subthalamic nucleus volume that became significant and negative after adjusting for brain size, suggesting that they contribute less to the g-factor than what is expected given their size. For mean thicknesses, regions include left and right transverse frontopolar gyri and sulci, the right lingual gyrus, right suborbital sulcus, the right superior frontal gyrus, the right cuneus gyrus, the right occipital superior gyrus, the right posterior-ventral part of the cingulate gyrus (isthmus), and left occipital pole and left posterior transverse collateral sulcus.

Across adjusted and unadjusted volumes, surface areas and thicknesses, and the different variants of our analyses, P-FIT regions were never overrepresented among the 20 or 30 regions that were most associated with g (Files S5–6) and they did not show larger associations with g than non-PFIT regions overall (Supplemental File S7).



Fig. 3. Meditating effect of regional cerebral measures on the g-factor PGS's effect on the g-factor with and without including childhood adversity. gfactor, general intelligence factor; PGS, polygenic scores. Coefficients correspond to direct effects. Fit of volume and surface area models: CFI = 1.00, SRMR=0.00, RMSEA=0.00. Fit of mean thickness model without childhood adversity: CFI=0.95, SRMR=0.05, RMSEA=0.15, and with CFI=0.95, SRMR=0.05, RMSEA=0.14. The PGS is adjusted for birth year and the first 40 principal components of the genotyped data. Cerebral measures and the g-factor are adjusted for sex, age, age², age by sex, age² by sex, and scanner site. By default, correlations between exogenous variables are fixed at their sample values in lavaan (Rosseel 2012). For simplicity, the error variance of each variable and correlations between mediators are not specified in the figure (Supplemental Tables D and E).

Do the regional measures that predict the g-factor independently from brain size still predict the g-factor when entered in the same model?

We then examined whether regions that significantly and positively or negatively predicted the g-factor after adjusting for brain size do so independently from each other. We found that there were 4 volumes, 3 surface areas, and 12 mean thicknesses that still significantly predicted the g-factor independently from each other and global brain size (Supplemental B9-11).

Do regional measures mediate the gPGS' effects on the g-factor? Volumes

Based on the previous analyses in Do the regional measures that predict the g-factor independently from brain size still predict the g-factor when entered in the same model? section, we included the right olfactory bulb, the left subcallosal gyrus, and the right mediodorsal medial magnocellular thalamic nuclei volumes in the regional and global volumetric mediation model. We did not include the third ventricle volume even if it was significant because we do not expect ventricular volumes to mediate genetic effects on intelligence.

The indirect path from the gPGS to the g-factor was significant for TBV and the right olfactory bulb volume after multiple comparison corrections and for the left subcallosal gyrus volume at the P < 0.05 threshold. TBV mediated 5.07% of the gPGS' effect on the g-factor, whereas the right olfactory bulb volume mediated 0.46% and left subcallosal gyrus volume 0.31% of the gPGS' effect on the g-factor (Fig. 3; Supplemental Table D3).

Surface areas

Based on the previous analyses in Do the regional measures that predict the g-factor independently from brain size still predict the g-factor when entered in the same model? section, we included TSA, the right orbital part of the inferior frontal gyrus surface area, the left subcallosal gyrus surface area, and the left anterior transverse collateral sulcus surface area in the regional and global surface area mediation model.

Total surface area mediated 3.99%, the left anterior transverse collateral sulcus surface area 0.41%, the left subcallosal gyrus 0.30%, and the right orbital part of the inferior frontal gyrus surface area 0.35% of the effect of the gPGS on the g-factor at the P < 0.05/4 threshold (Fig. 3; Supplemental Table D4).

Mean thicknesses

Based on the previous analyses in Do the regional measures that predict the g-factor independently from brain size still predict the g-factor when entered in the same model? section, we included total MCT, the right anterior part of the cingulate gyrus and sulcus, the left anterior part of the cingulate gyrus and sulcus, the left planum polare of the superior temporal gyrus, the left inferior segment of the circular sulcus of the insula, the right pericallosal sulcus, the left short insular gyrus, the right postcentral sulcus, the right superior segment of circular insula sulcus, the right superior temporal sulcus, the right precentral gyrus, the left subcallosal gyrus, and the left pericallosal sulcus mean thicknesses in the regional and global mean thickness mediation model.

Total MCT did not mediate the effects of the gPGS or childhood adversity on the g-factor. The right anterior part of the cingulate gyrus and sulcus mean thickness mediated 0.27% of the effect of the gPGS on the g-factor at P < 0.05. The left anterior part of the cingulate gyrus mediated 0.14%, the left short insular gyrus mean thickness 0.18%, the left planum polare of the superior temporal gyrus mean thickness 0.16%, the right pericallosal sulcus mean thickness mediated 0.12%, the left inferior segment of the circular sulcus of the insula mean thickness mediated 0.18%, and the right pericallosal sulcus mediated 0.27% of the effect of the gPGS on the g-factor at P < 0.05 (Fig. 3; Supplemental Table D5).

Do regional measures mediate the gPGS' and childhood adversity's effects on the g-factor? Volumes

In the volumetric mediation models, TBV mediated 4.15% of the gPGS' effect on the g-factor and the right olfactory bulb volume mediated 0.37% of the gPGS' effect on the g-factor at P < 0.05/4, whereas the right mediodorsal medial magnocellular thalamic nuclei volume mediated 0.32% of the gPGS' effect on the g-factor at P < 0.05. Therefore, the left subcallosal gyrus volume was no longer significant when adding childhood adversity in the mediation model and the indirect path through the right mediodorsal medial magnocellular thalamic nuclei became significant at P < 0.05. TBV mediated 1.41% of childhood adversity's effect on the g-factor, whereas the right olfactory bulb volume mediated 0.14% of childhood adversity's effect on the g-factor at P < 0.05 (Fig. 3; Supplemental Table E3).

Surface areas

Total surface area mediated 3.37%, the left anterior transverse collateral sulcus 0.34%, and the right orbital part of the inferior frontal gyrus surface area 0.29% of the effect of the gPGS on the g-factor at the P < 0.05/4 threshold (Supplemental Table D4). Total surface area mediated 1.11% and the left anterior transverse collateral sulcus surface area 0.16% of childhood adversity's effect on the g-factor (Fig. 3; Supplemental Table E4).

Mean thicknesses

Total MCT did not mediate the effects of the gPGS or childhood adversity on the g-factor. The right anterior part of the cingulate gyrus and sulcus mean thickness still mediated 0.27% of the effect of the gPGS on the g-factor at P < 0.05. The right pericallosal sulcus mean thickness mediated 0.12%, the left short insular gyrus mean thickness 13%, and the left planum polare of the superior temporal gyrus mean thickness 0.15% of the effect of the gPGS on the g-factor at P < 0.05. However, the left inferior segment of the circular sulcus of the insula mean thickness and the left anterior part of the cingulate gyrus and sulcus mean thickness no longer mediated the gPGS' effect on the g-factor: Instead, a new region mediated the gPGS' effect on the g-factor: the mean thickness of left subcallosal gyrus (0.17%), which also mediated the effect of childhood adversity on the g-factor (0.10%; Fig. 3; Supplemental Table C5).

Discussion

This paper examined the neuroanatomical measures (e.g. brain volumes) mediating the effect of genes and childhood adversity on intelligence. Both genetic and environmental components were associated with intelligence and exerted their influence on the brain at the macro-structural level. Our first analyses on the neural correlates of intelligence revealed that (i) some regions are correlated with intelligence beyond their association with global brain size and their associations with other brain regions and that (ii), while functional studies may support the P-FIT theory, our structural findings do not support the P-FIT. When examining whether global measures and regions that uniquely contribute to the g-factor mediate the gPGS' and childhood adversity's effects on the g-factor, we found that (i) both environmental and genetic effects were uniquely associated with intelligence, (ii) the effect of genes and childhood adversity were mostly mediated by global effects, and (iii) the effects of childhood adversity and genes on the g-factor were mediated by similar regions. However, TBV, the largest mediator, mediated about 2.5% and 10% of childhood adversity and the gPGS' effects on the g-factor, respectively, suggesting that mediation by neuroanatomical measures only explains a small fraction of the total effect of the gPGS and childhood adversity on the g-factor.

Most associations between the g-factor and volumes (78%) or surface areas (88%) disappeared after adjusting for global brain size, suggesting that the majority of volumes and surfaces contribute to intelligence through global cerebral effects. In contrast, only 4% of mean thicknesses were no longer significant after adjusting for global brain size and 26% of mean thicknesses still predicted the g-factor after adjusting for total MCT. Therefore, mean thicknesses appear to influence the g-factor through region-specific effects rather than global effects. This can be explained by the small associations between regional mean thicknesses and total MCT (mean $\beta = 0.03$) compared to those of regional volumes with TBV (mean $\beta = 0.30$) as well as the small association between total MCT and the g-factor ($\beta = 0.04$). Adjusting for total MCT thus captures little variance between regional mean cortical thicknesses and the g-factor.

Regions that contributed the most to the g-factor not only corresponded to regions that significantly predicted the g-factor after adjusting for brain size but also maintained the direction of their effect with and without adjusting for global measures. We found that cerebellar volumes, subcortical nuclei volumes, a few cortical volumes, frontal surface areas, and temporal mean thicknesses contributed more positively to the gfactor than what was expected given their size. In contrast, a few distributed mean thicknesses and ventricular volumes contributed more negatively to the g-factor than what was expected given their size. Negative and positive associations between cortical thickness and intelligence have been reported across the cortex (Karama et al. 2014) and are thought to depend on the measure of the intelligence (Goriounova and Mansvelder 2019): Greater crystallized intelligence is associated with cortical thinning, whereas fluid intelligence does not appear to be related with cortical thickness (Tadayon et al. 2020). Studies looking at age-related changes in performance on the cognitive tests of the UK Biobank found that performance on the verbal numerical UK Biobank test (also known as the fluid intelligence test) does not decrease as expected with age (Hagenaars et al. 2016). Instead, the performance stagnates as would be expected with crystallized intelligence (Cavanaugh and Blanchard-Fields 2018). Therefore, our g-factor measure, on which the fluid intelligence test loads highly (0.62), likely captures both crystallized and fluid intelligence and may explain why we find distributed positive and negative associations between mean thicknesses and intelligence. Future studies should explore the associations between brain regions and subdomains of intelligence to obtain a better understanding of the associations between general intelligence, cognitive abilities, and the cortex (Jung and Haier 2007).

When approximately matching the Destrieux segmentations to the P-FIT Brodmann areas, the proportion of P-FIT and non-P-FIT regions associated with the g-factor was generally similar across cortical volumes, surface areas, and mean thicknesses and the magnitude of the association was not larger for P-FIT regions. Therefore, although we report associations with frontal and temporal-parietal regions that are concordant with the P-FIT, our findings do not support this theory with regard to volumes, thicknesses, and surface areas. However, the P-FIT may nonetheless accurately predict functional brain activations associated with general intelligence (Haier and Jung 2018; Gur et al. 2021).

Concerning other regions of interest, we did not replicate the significant associations between the g-factor and the whole hippocampal or thalamic volumes after adjusting for brain size previously reported by a UK Biobank study (Cox et al. 2019). Instead, we report associations between the g-factor and subcortical subsegmentations as well as cerebellar subsegmentations. For instance, we find a positive association between the g-factor and the right mediodorsal thalamic nucleus volume, a region known to critically contribute to cognitive functions (Ouhaz et al. 2018), and positive associations between the g-factor and most of the crus lobules of the cerebellum, which are functionally connected to regions of the default mode network (Buckner et al. 2011), a network of higher-level cognition (Smallwood et al. 2021).

We find that regional measures independently explained a small portion of the effect of the gPGS on the g-factor and that the gPGS' effect on the g-factor is mediated by several cortical surface areas, volumes, and mean thickness. Although we used different segmentations and samples from Lett et al. (2020), we similarly find that the gPGS' effect on the g-factor is mediated by the anterior cingulate cortex, prefrontal, insular, medial temporal, and inferior parietal mean thicknesses and surface areas. However, we find that our regional measures mediate a smaller percentage of the gPGS' effect on the g-factor (around 0.30% instead of 0.75%), which may be due to the difference in age between cohorts or to their larger segmentations. We additionally report volumes that mediate the gPGS' effect, and most notably the right mediodorsal medial magnocellular thalamic nuclei, a region thought to be implicated in executive functions (e.g. cognitive control and decision-making; Ouhaz et al. 2018).

When adding childhood adversity to the model, the percentage mediated by regional and global effects decreased to various extents across regions. For instance, adding childhood adversity to the mediation models did not impact the percentage mediated by the mean thickness of the right anterior part of the cingulate gyrus and sulcus. However, the surface area and volume of the left subcallosal gyrus and the mean thicknesses of the left inferior segment of the circular sulcus of the insula and the left anterior part of the cingulate gyrus and sulcus no longer mediated the gPGS' effect on the g-factor when adding childhood adversity to the model. Considering that childhood adversity significantly and negatively predicted these regions, the association between these non-longer mediating regions may be due to the correlation between childhood adversity and the gPGS: Part of the variance previously attributed to the gPGS may have shifted from the gPGS to childhood adversity. Finally, when adding childhood adversity to the mean thickness mediation model, a new region mediated both the gPGS effect and childhood adversity's effect on the g-factor: the left subcallosal gyrus mean thickness. The latter highlights the importance of including environmental measures to better understand the complex relationship between environmental and genetic effects on general intelligence.

Although we find that specific regions mediate the g-factor and childhood adversity's effects independently from global brain size and regional associations, the mediation of global brain size was 10–20 times larger than the mediation of specific regions when examining volumes and surface areas. TBV explained 2.3% and the regional volumes included in the mediation models explained 0.3% of the variance in the g-factor, whereas TSA explained 1.8% and the regional surface areas included in the mediation models explained 0.2% of the variance in the g-factor. These findings are consistent with previous studies suggesting that general intelligence may be more related to global than regionspecific differences in the gray matter volume (Hilger et al. 2020) and that adding regional effects on the g-factor does not substantially predict more variance in the g-factor than TBV alone (Cox et al. 2019). However, TBV only mediated 7.04% of the gPGS' effect on the g-factor and 2.50% of childhood adversity's effect on the g-factor, leaving 93% of the gPGS' effect on the g-factor, and 97.5% of childhood adversity's effect on the g-factor to be explained by other cerebral measures. Therefore, future research should include additional cerebral measures, such as microstructural properties, white matter measures, dynamic connectivity, or resting-state or task-based functional activation, to better understand the extent to which cerebral measures mediate the environmental and genetic factors on the g-factor.

The present study is limited in its ability to make causal claims regarding the mediation of neuroanatomical structures on intelligence. Although the mediation model assumes that brain measures cause intelligence scores, intelligence scores may influence the investigated brain measures because such measures are to some extent plastic, and both are measured around the same time in older adulthood. Therefore, whereas the effects of the gPGS and childhood adversity on brain measures and intelligence during older adulthood may be causal, associations between brain measures and intelligence scores are correlational in nature. Future studies may be able to address this caveat by using brain measures that are taken after measures of childhood adversity but before measures of intelligence.

The paper is also limited in its ability to generalize to all UK Biobank participants: Individuals with neuroimaging data are different from UK Biobank participants without neuroimaging data (Lyall et al. 2022) and the UK population (Fry et al. 2017). Our analyses were restrained to individuals of British ancestry, suggesting that further research is needed to examine whether genetic factors on intelligence are mediated by the same cerebral regions and to the same degree across ancestries. The PGS also only predicted 7.9% of the variance in the g-factor, suggesting that additional regions may be found when using a more predictive PGS. Finally, not including the most important environmental predictor of intelligence, parental or childhood socioeconomic status (SES; Flensborg-Madsen et al. 2020; Flensborg-Madsen and Mortensen 2017), is a major limitation of our study. Since childhood SES was not available in the UK Biobank, we focused on childhood adversity, which is strongly associated with intelligence (but less so than parental SES; McGuire and Jackson 2020). Although we could have used adult SES, as done in previous studies (e.g. Kweon et al. 2022) to serve as a proxy of childhood SES, the bidirectional influences between adult SES and intelligence would have compromised the interpretation of the mediation model.

The present paper provides the first large-scale study examining the neuroanatomical measures mediating genetic (gPGS) and early environmental (childhood adversity) effects on intelligence (g-factor). We replicate and extend previous findings and highlight the importance of adding environmental data to better understand how genetic and environmental factors act on the brain to influence general intelligence. In light of the strong evidence for genetic and environmental factors contributing to individual differences in intelligence (for review Deary et al. 2021; Harden 2021), we urge future studies to simultaneously investigate the genetic, environmental, and cerebral effects on intelligence by examining a variety of cerebral properties, from the macro to the micro, to understand discrepancies in intelligence (for review Deary et al. 2021) and, in turn, later health, educational, and social outcomes (Schmidt and Hunter 2004; Strenze 2007; Calvin et al. 2017; Twig et al. 2018).

Acknowledgments

We thank the UK Biobank Resource for making this research possible.

Supplementary material

Supplementary material is available at Cerebral Cortex online.

Funding

This work received support under the program "Investissements d'Avenir" launched by the French Government and implemented by l'Agence Nationale de la recherche (ANR) with the references ANR-17-EURE-0017 and ANR-10-IDEX-0001-02 PSL. A CC-BY public copyright license has been applied by the authors to the present document and will be applied to all subsequent versions up to the Author Accepted Manuscript arising from this submission, in accordance with the grant's open access conditions. This research has been conducted using the UK Biobank Resource.

Conflict of interest statement: On behalf of all authors, the corresponding author states that there is no conflict of interest.

Data/code availability

This research has been conducted using data from UK Biobank (http://www.ukbiobank.ac.uk/). Restrictions apply to the availability of these data, which were used under license for this study: application 46007. Preregistration, supplemental tables, and supplemental files, and the code are available here: https:// osf.io/ec97u/?view_only=4b366bd7ed2442a1a9f64bfcc2fe0946.

References

- Alfaro-Almagro F, Jenkinson M, Bangerter NK, Andersson JLR, Griffanti L, Douaud G, Sotiropoulos SN, Jbabdi S, Hernandez-Fernandez M, Vallee E, et al. Image processing and quality control for the first 10,000 brain imaging datasets from UK Biobank. *NeuroImage*. 2018:166:400–424.
- Basten U, Hilger K, Fiebach CJ. Where smart brains are different: a quantitative meta-analysis of functional and structural brain imaging studies on intelligence. Intelligence. 2015:51:10–27.
- Bates TC, Maher BS, Medland SE, McAloney K, Wright MJ, Hansell NK, Kendler KS, Martin NG, Gillespie NA. The nature of nurture: using a virtual-parent design to test parenting effects on children's educational attainment in genotyped families. Twin Res Human Genetics. 2018:21(2):73–83.
- Buckner RL, Krienen FM, Castellanos A, Diaz JC, Yeo BTT. The organization of the human cerebellum estimated by intrinsic functional connectivity. J Neurophysiol. 2011:106(5):2322–2345.
- Calvin CM, Batty GD, Der G, Brett CE, Taylor A, Pattie A, Čukić I, Deary IJ. Childhood intelligence in relation to major causes of death in 68 year follow-up: prospective population study. *BMJ*. 2017:357.

- Cavanaugh JC, Blanchard-Fields F. Adult development and aging. Cengage Learning; 2018
- Colom R, Karama S, Jung RE, Haier RJ. Human intelligence and brain networks. Dialogues Clin Neurosci. 2010:12(4):489–501.
- R Core Team. (2022). R: a language and environment for statistical computing. R Foundation for Statistical Computing. https://www.R-project.org/
- Cox SR, Ritchie SJ, Tucker-Drob EM, Liewald DC, Hagenaars SP, Davies G, Wardlaw JM, Gale CR, Bastin ME, Deary IJ. Ageing and brain white matter structure in 3,513 UK Biobank participants. *Nat Commun.* 2016:7:13629.
- Cox SR, Ritchie SJ, Fawns-Ritchie C, Tucker-Drob EM, Deary IJ. Structural brain imaging correlates of general intelligence in UK Biobank. Intelligence. 2019:76:101376.
- Deary IJ, Penke L, Johnson W. The neuroscience of human intelligence differences. Nat Rev Neurosci. 2010:11(3):Article 3.
- Deary IJ, Cox SR, Hill WD. Genetic variation, brain, and intelligence differences. Mol Psychiatry. 2021:27:335–353.
- Dye H. The impact and long-term effects of childhood trauma. J Hum Behav Soc Environ. 2018:28(3):381–392.
- Enlow MB, Egeland B, Blood EA, Wright RO, Wright RJ. Interpersonal trauma exposure and cognitive development in children to age 8 years: a longitudinal study. J Epidemiol Community Health. 2012:66(11):1005–1010.
- Flensborg-Madsen T, Mortensen EL. Birth weight and intelligence in young adulthood and midlife. *Pediatrics*. 2017:139(6):e20163161.
- Flensborg-Madsen T, Eriksen H-LF, Mortensen EL. Early life predictors of intelligence in young adulthood and middle age. PLoS One. 2020:15(1):e0228144.
- Fry A, Littlejohns TJ, Sudlow C, Doherty N, Adamska L, Sprosen T, Collins R, Allen NE. Comparison of sociodemographic and healthrelated characteristics of UK biobank participants with those of the general population. Am J Epidemiol. 2017:186(9):1026–1034.
- Genç E, Metzen D, Fraenz C, Schlüter C, Voelkle MC, Arning L, Streit F, Nguyen HP, Güntürkün O, Ocklenburg S, et al. Structural architecture and brain network efficiency links polygenic scores to intelligence. *bioRxiv*. 2022:2022.03.22.485284.
- Gignac GE, Bates TC. Brain volume and intelligence: the moderating role of intelligence measurement quality. *Intelligence*. 2017:64: 18–29.
- Goriounova NA, Mansvelder HD. Genes, cells and brain areas of intelligence. Front Hum Neurosci. 2019:13.
- Gur RC, Butler ER, Moore TM, Rosen AFG, Ruparel K, Satterthwaite TD, Roalf DR, Gennatas ED, Bilker WB, Shinohara RT, et al. Structural and functional brain parameters related to cognitive performance across development: replication and extension of the parieto-frontal integration theory in a single sample. *Cereb* Cortex. 2021:31(3):1444–1463.
- Hagenaars SP, Harris SE, Davies G, Hill WD, Liewald DCM, Ritchie SJ, Marioni RE, Fawns-Ritchie C, Cullen B, Malik R, et al. Shared genetic aetiology between cognitive functions and physical and mental health in UK Biobank (N=112 151) and 24 GWAS consortia. *Mol Psychiatry*. 2016:21(11):Article 11.
- Haier RJ, Jung RE. The parieto-frontal integration theory: assessing intelligence from brain images. In: DP Flanagan, EM McDonough (eds.), Contemporary intellectual assessment: theories, tests, and issues. NY: The Guilford Press; 2018:pp. 219–224.
- Harden KP. "Reports of My Death Were Greatly Exaggerated": behavior genetics in the postgenomic era. *Annu Rev Psychol*. 2021:72(1): 37–60.
- Haworth CMA, Wright MJ, Luciano M, Martin NG, de Geus EJC, van Beijsterveldt CEM, Bartels M, Posthuma D, Boomsma DI, Davis OSP, et al. The heritability of general cognitive ability

increases linearly from childhood to young adulthood. Mol Psychiatry. 2010:15(11):1112–1120.

- Hilger K, Winter NR, Leenings R, Sassenhagen J, Hahn T, Basten U, Fiebach CJ. Predicting intelligence from brain gray matter volume. Brain Struct Funct. 2020:225(7):2111–2129.
- Hill WD, Marioni RE, Maghzian O, Ritchie SJ, Hagenaars SP, McIntosh AM, Gale CR, Davies G, Deary IJ. A combined analysis of genetically correlated traits identifies 187 loci and a role for neurogenesis and myelination in intelligence. *Mol Psychiatry*. 2019:24(2): Article 2.
- Hu L, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. Structural Equation Modeling: A Multidisciplinary Journal. 1999:6(1): 1–55.
- Johnson MR, Shkura K, Langley SR, Delahaye-Duriez A, Srivastava P, Hill WD, Rackham OJL, Davies G, Harris SE, Moreno-Moral A, et al. Systems genetics identifies a convergent gene network for cognition and neurodevelopmental disease. *Nat Neurosci.* 2016:19(2): 223–232.
- Jorgensen TD, Pornprasertmanit S, Schoemann AM, Rosseel Y. sem-Tools: Useful tools for structural equation modeling. *R package version* 0.5–6. 2022. Retrieved from https://CRAN.R-project.org/ package=semTools.
- Jung RE, Haier RJ. The parieto-frontal integration theory (P-FIT) of intelligence: converging neuroimaging evidence. *Behav Brain Sci.* 2007:30(2):135–154 discussion 154-187.
- Karama, S., Bastin, M. E., Murray, C., Royle, N. A., Penke, L., Muñoz Maniega, S., Gow, A. J., Corley, J., Valdés Hernández, M., Lewis, J. D., et al. Childhood cognitive ability accounts for associations between cognitive ability and brain cortical thickness in old age. *Mol Psychiatry*. 2014:19(5):Article 5:555–559.
- Kweon H, Aydogan G, Dagher A, Bzdok D, Ruff CC, Nave G, Farah MJ, Koellinger PD. Human brain anatomy reflects separable genetic and environmental components of socioeconomic status. Sci Adv. 2022:8(20):eabm2923.
- Lee JJ, Wedow R, Okbay A, Kong E, Maghzian O, Zacher M, Nguyen-Viet TA, Bowers P, Sidorenko J, Linnér RK, et al. Gene discovery and polygenic prediction from a genome-wide association study of educational attainment in 1.1 million individuals. *Nat Genet*. 2018:50(8):1112–1121.
- Lett TA, Vogel BO, Ripke S, Wackerhagen C, Erk S, Awasthi S, Trubetskoy V, Brandl EJ, Mohnke S, Veer IM, et al. Cortical surfaces mediate the relationship between polygenic scores for intelligence and general intelligence. *Cereb Cortex*. 2020:30(4):2708–2719.
- Lloyd-Jones LR, Zeng J, Sidorenko J, Yengo L, Moser G, Kemper KE, Wang H, Zheng Z, Magi R, Esko T, et al. Improved polygenic prediction by Bayesian multiple regression on summary statistics. Nat Commun. 2019:10(1):Article 1. https://doi.org/10.1038/ s41467-019-12653-0.
- Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. Nat Rev Neurosci. 2009:10(6):Article 6.
- Lyall DM, Quinn T, Lyall LM, Ward J, Anderson JJ, Smith DJ, Stewart W, Strawbridge RJ, Bailey MES, Cullen B. Quantifying bias in psychological and physical health in the UK Biobank imaging sub-sample. *Brain Commun.* 2022:4(3):fcac119.
- McGuire A, Jackson Y. The role of trauma type and age in the relation between trauma exposure and intelligence. *Child Maltreatment*. 2020:25(2):192–202.

- Miller KL, Alfaro-Almagro F, Bangerter NK, Thomas DL, Yacoub E, Xu J, Bartsch AJ, Jbabdi S, Sotiropoulos SN, Andersson JLR, et al. Multimodal population brain imaging in the UK Biobank prospective epidemiological study. Nat Neurosci. 2016:19(11):1523–1536.
- Morris TT, Davies NM, Davey Smith G. Can education be personalised using pupils' genetic data? *elife*. 2020:9:e49962.
- Ouhaz Z, Fleming H, Mitchell AS. Cognitive functions and neurodevelopmental disorders involving the prefrontal cortex and mediodorsal thalamus. Front Neurosci. 2018:12:33. https://www. frontiersin.org/article/10.3389/fnins.2018.00033.
- Pietschnig J, Penke L, Wicherts JM, Zeiler M, Voracek M. Meta-analysis of associations between human brain volume and intelligence differences: How strong are they and what do they mean? Neurosci Biobehav Rev. 2015:57:411–432.
- Polderman TJC, Benyamin B, de Leeuw CA, Sullivan PF, van Bochoven A, Visscher PM, Posthuma D. Meta-analysis of the heritability of human traits based on fifty years of twin studies. *Nat Genet.* 2015:47(7):Article 7.
- Rimfeld K, Malanchini M, Krapohl E, Hannigan LJ, Dale PS, Plomin R. The stability of educational achievement across school years is largely explained by genetic factors. NPJ Sci Learn. 2018:3(1):1–10.
- Rosseel Y. lavaan: an R Package for structural equation modeling. J Stat Softw. 2012:48(2):1–36.
- Saarentaus EC, Havulinna AS, Mars N, Ahola-Olli A, Kiiskinen TTJ, Partanen J, Ruotsalainen S, Kurki M, Urpa LM, Chen L, et al. Polygenic burden has broader impact on health, cognition, and socioeconomic outcomes than most rare and high-risk copy number variants. Mol Psychiatry. 2021:26(9):4884–4895.
- Savage JE, Jansen PR, Stringer S, Watanabe K, Bryois J, de Leeuw CA, Nagel M, Awasthi S, Barr PB, Coleman JRI, et al. Genome-wide association meta-analysis in 269,867 individuals identifies new genetic and functional links to intelligence. *Nat Genet*. 2018:50(7): 912–919.
- Schmidt FL, Hunter J. General mental ability in the world of work: occupational attainment and job performance. J Pers Soc Psychol. 2004:86(1):162–173.
- Smallwood J, Bernhardt BC, Leech R, Bzdok D, Jefferies E, Margulies DS. The default mode network in cognition: a topographical perspective. Nat Rev Neurosci. 2021:22(8):Article 8.
- Strenze T. Intelligence and socioeconomic success: a meta-analytic review of longitudinal research. Intelligence. 2007:35(5):401–426.
- Tadayon E, Pascual-Leone A, Santarnecchi E. Differential contribution of cortical thickness, surface area, and gyrification to fluid and crystallized intelligence. Cereb Cortex (New York, NY). 2020:30(1):215–225.
- Twig G, Tirosh A, Derazne E, Haklai Z, Goldberger N, Afek A, Gerstein HC, Kark JD, Cukierman-Yaffe T. Cognitive function in adolescence and the risk for premature diabetes and cardiovascular mortality in adulthood. Cardiovasc Diabetol. 2018:17(1):154.
- von Stumm S, Smith-Woolley E, Ayorech Z, McMillan A, Rimfeld K, Dale PS, Plomin R. Predicting educational achievement from genomic measures and socioeconomic status. *Dev Sci.* 2020:23(3):e12925.
- Williams CM, Peyre H, Toro R, Ramus F. Neuroanatomical norms in the UK Biobank: the impact of allometric scaling, sex, and age. *Hum Brain Mapp.* 2021:42(14):4623–4642.
- Williams CM, Labouret G, Wolfram T, Peyre H, Ramus F. A general cognitive ability factor for the UK Biobank. Behav Genet. 2022. https://doi.org/10.1007/s10519-022-10127-6.